1. An optically active compound of the formula:

5

where the R_2 and R_3 groups are methyl, another lower alkyl group or an aryl or biaryl unit while the R_1 groups independently each are a hydroxyl, alkoxyl, aryloxy, or arylalkoxy group, the R groups each represent a group as follows:

$$A_1 - [-Z-]_q - A_2 -$$

10

where A_1 is an aromatic group, an acyclic aliphatic group, or an alicyclic group, and A_1 can be a substituted or unsubstituted group, Z is a group selected from -O-, -OCO-, or -S-, and the coefficient q is 0 or 1. Z may also be $(CH_2)_nO$ where the coefficient n is 0 to 5 and the coefficient q is 1. A_2 is a bivalent radical of a naphthalene group, and the cyclic structure of A_2 , or A_1 if it is cyclic, optionally can be heterocyclic, such as by replacement of one or more CH member(s) of the ring structure with N, O and/or S.

15

2. The optically active compound of claim 1, where each R substituent is independently selected as:

20

where R₄ represents a group as follows:

$$Y - [-X-]_n - [-Z-]_q$$

KD-01-004PCT

where n is an integer value of 0 or 1 or more, X is -CH=CH-CH₂-, or -(CH₂)_m- where m is an integer value of 1, 2, 3, or more, Y is a radical of an aromatic hydrocarbon, an acyclic aliphatic hydrocarbon, or an alicyclic hydrocarbon, and Y can be a substituted or unsubstituted group, and Z and q have the same respective meanings as defined in claim 1.

5

- 3. The optically active compound of claim 2, where R₄ is an aryloxy radical, an arylalkoxy radical, an arylalkyleneoxy, or an arylalkenyleneoxy radical.
- 4. (4R,5R)-2,2-dimethyl-α,α,α',α'-tetrakis[6-(benzyloxy)naphth-2-yl]-1,3-dioxolane4.5-dimethanol.
 - 5. A liquid crystalline mixture, comprising: a liquid-crystalline base having liquid crystalline properties; at least one optically active compound of the formula as described in any of one of claims 1-4.

15

20

6. The liquid crystalline mixture according to claim 5, further including an achiral non-liquid crystalline compound.

7. The liquid crystalline mixture according to claim 6, wherein the achiral non-liquid crystalline compound comprises R^1 -C \equiv N, where R^1 represents an aliphatic group.

25

30

8. The liquid crystalline mixture according to claim 7, wherein R¹-C≡N comprises an alkylnitrile.

9. The liquid crystalline mixture according to claim 7, wherein R¹-C≡N comprises

undecanenitrile.

10. An electro-optical cell comprising a layer including a liquid crystalline mixture as described in any one of claims 5-9 sandwiched between two substrate means, and means for applying an electric potential to the substrate means.

10

15

20

- 11. A light modulating apparatus comprising an electro-optical cell according to claim 10.
- 5 12. The light modulating apparatus according to claim 11, wherein the light modulating apparatus comprises a cholesteric display.
 - 13. An electro-optical cell comprising:
 - a layer comprising:

about 2-6 wt% achiral material.

at least 70 weight percent (wt%) nematic host mixture; and at least about 2 wt% (4R,5R)-2,2-dimethyl-α,α,α',α'-tetrakis[6-

(benzyloxy)naphth-2-yl]-1,3-dioxolane-4,5-dimethanol;

first and second substrates disposed above and below, respectively, the layer; and first and second conductors physically coupled to the first and second substrates, respectively, which permit an electrical potential to be applied across the layer.

- 14. The electro-optical cell as recited in claim 13, wherein the layer further comprises
- 15. The electro-optical cell as recited in claim 13, wherein the layer further comprises a chiral material different from (4R,5R)-2,2-dimethyl-α,α,α',α'-tetrakis[6-(benzyloxy)naphth-2-yl]-1,3-dioxolane-4,5-dimethanol and having an opposite twist sense.
- 16. A light modulating apparatus comprising an electro-optical cell according to any 25 one of claims 13-15.
 - 17. The light modulating apparatus according to claim 16, wherein the lightmodulating apparatus comprises a cholesteric display having a temperature independent reflective wavelength.